

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 March 2003 (27.03.2003)

PCT

(10) International Publication Number
WO 03/024386 A2

(51) International Patent Classification⁷: **A61K**
(21) International Application Number: PCT/IB02/04236
(22) International Filing Date:
20 September 2002 (20.09.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/323,315 20 September 2001 (20.09.2001) US

(71) Applicant (for all designated States except US): **AB SCIENCE** [FR/FR]; 3, avenue Georges V, F-75008 Paris (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MOUSSY, Alain** [FR/FR]; 22 bis, passage Dauphine, F-75006 Paris (FR).
KINET, Jean-Pierre [FR/US]; 3 Hunt Road, Lexington, MA 02421 (US).

(74) Agents: **MARTIN, Jean-Jacques** et al.; Cabinet Regimbeau, 20, rue de Chazelles, F-75847 Paris Cedex 17 (FR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

Published:

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF POTENT, SELECTIVE AND NON TOXIC C-KIT INHIBITORS FOR TREATING INTERSTITIAL CYSTITIS

(57) Abstract: The present invention relates to a method for treating interstitial cystitis, comprising administering a tyrosine kinase inhibitor to a human in need of such treatment, more particularly a non-toxic, potent and selective c-kit inhibitor, wherein said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.



WO 03/024386 A2

Use of potent, selective and non toxic c-kit inhibitors
for treating interstitial cystitis

- 5 The present invention relates to a method for treating interstitial cystitis, comprising administering a tyrosine kinase inhibitor to a human in need of such treatment, more particularly a non toxic, potent and selective c-kit inhibitor, wherein said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.
- 10 Interstitial cystitis (IC) is a chronic inflammation of the bladder wall resulting in tissue damage, especially at the interstices between the cells in the lining of the bladder. IC affects up to 700,000 women in the United States. The symptoms include pain, urgency and frequency of urination and cystoscopic abnormalities including hemorrhages, Oravisto, K. J. (1975) Ann. Chir. Gynaecol. Fenn. 64: 75. As a result, quality of life
- 15 scores in IC patients are very low. Moreover, as of today, none of the proposed medications provide a cure.

The hypothesized causes of IC include infectious, lymphovascular obstruction and neurogenic, endocrinologic, psychoneurotic, inflammatory and autoimmune pathologies.

- 20 The fact that lymphocytes infiltrate into the bladder wall of patients is of particular interest. Bladder inflammation is also illustrated by a significant increase both in mast cell number and size. Furthermore, Histopathological studies have demonstrated that mast cells in the bladder walls of IC patients are degranulated.

- 25 Mast cells (MC) are tissue elements derived from a particular subset of hematopoietic stem cells that express CD34, c-kit and CD13 antigens (Kirshenbaum et al, Blood. 94: 2333-2342, 1999 and Ishizaka et al, Curr Opin Immunol. 5: 937-43, 1993). Immature

MC progenitors circulate in the bloodstream and differentiate in tissues. These differentiation and proliferation processes are under the influence of cytokines, one of utmost importance being Stem Cell Factor (SCF), also termed Kit ligand (KL), Steel factor (SL) or Mast Cell Growth Factor (MCGF). SCF receptor is encoded by the protooncogene c-kit, that belongs to type III receptor tyrosine kinase subfamily (Boissan and Arock, J Leukoc Biol. 67: 135-48, 2000). This receptor is also expressed on others hematopoietic or non hematopoietic cells. Ligation of c-kit receptor by SCF induces its dimerization followed by its transphosphorylation, leading to the recruitment and activation of various intracytoplasmic substrates. These activated substrates induce multiple intracellular signaling pathways responsible for cell proliferation and activation (Boissan and Arock, 2000). Mast cells are characterized by their heterogeneity, not only regarding tissue location and structure but also at the functional and histochemical levels (Aldenborg and Enerback., Histochem. J. 26: 587-96, 1994 ; Bradding et al. J Immunol. 155: 297-307, 1995 ; Irani et al, J Immunol. 147: 247-53, 1991 ; Miller et al, Curr Opin Immunol. 1: 637-42, 1989 and Welle et al, J Leukoc Biol. 61: 233-45, 1997).

Here, it is proposed that mast cells are directly or indirectly implicated in the inflammation observed in IC and could lead to the destruction of the interstices between cells of the bladder wall.

20

Sant GR et al, Urol Clin North Am 1994 Feb;21(1):41-53 have shown that mast cells found in the bladder contain many granules, each of which can secrete many vasoactive and nociceptive molecules. In addition, according to Saban R et al , Physiol Genomics 2001 Aug 8, bladder inflammation does not occur in mast cell-deficient (Kit(W)/Kit(W-v)), whereas inflammation is observed upon stimuli in wild mice.

25

In connection with the present invention, it is postulated that activated mast cells secrete a number of cytokines and proteases that damage bladder mucosa while also attracting other inflammatory cells such as T lymphocytes and macrophages, which further participate in the inflammation and destruction process.

5 Indeed, upon stimuli, mast cells produce a large variety of mediators categorized into three groups:

- preformed granule-associated mediators (histamine, proteoglycans, and neutral proteases),
- lipid-derived mediators (prostaglandins, thromboxanes and leucotrienes),
- 10 - and various cytokines (IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-8, TNF- α , GM-CSF, MIP-1a, MIP-1b and IFN- γ).

More specifically, an SCF/IL-6-driven mast cells response has been found in IC, which shows that mast cell play a crucial role in the genesis and development of interstitial cystitis.

15

Therefore, the invention provides a new therapeutic strategy aimed at the use of c-kit specific kinase inhibitors to inhibit mast cell proliferation, survival and activation. A new route for treating interstitial cystitis is provided, which consists of destroying mast cells that are involved in the destruction of bladder muscosa.

20

It has been found that tyrosine kinase inhibitors and more particularly c-kit inhibitors that are unable to promote death of IL-3 dependent cells cultured in presence of IL-3 are especially suited to reach this goal.

25

Description

The present invention relates to a method for treating interstitial cystitis comprising administering a tyrosine kinase inhibitor to a human in need of such treatment, wherein
5 said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.


Tyrosine kinase inhibitors are selected for example from bis monocyclic, bicyclic or heterocyclic aryl compounds (WO 92/20642), vinylene-azaindole derivatives (WO
10 94/14808) and 1-cyclopropyl-4-pyridyl-quinolones (US 5,330,992), Styryl compounds (US 5,217,999), styryl-substituted pyridyl compounds (US 5,302,606), seleoindoles and selenides (WO 94/03427), tricyclic polyhydroxylic compounds (WO 92/21660) and benzylphosphonic acid compounds (WO 91/15495), pyrimidine derivatives (US 5,521,184 and WO 99/03854), indolinone derivatives and pyrrol-substituted indolinones
15 (US 5,792,783, EP 934 931, US 5,834,504, US 5,883,116, US 5,883,113, US 5, 886,020, WO 96/40116 and WO 00/38519), as well as bis monocyclic, bicyclic aryl and heteroaryl compounds (EP 584 222, US 5,656,643 and WO 92/20642), quinazoline derivatives (EP 602 851, EP 520 722, US 3,772,295 and US 4,343,940) and aryl and heteroaryl quinazoline (US 5,721,237, US 5,714,493, US 5,710,158 and WO 95/15758).

20

Preferably, said tyrosine kinase inhibitors are non-toxic, selective and potent c-kit inhibitors. Such inhibitors can be selected from the group consisting of indolinones, pyrimidine derivatives, pyrrolopyrimidine derivatives, quinazoline derivatives, quinoxaline derivatives, pyrazoles derivatives, bis monocyclic, bicyclic or heterocyclic
25 aryl compounds, vinylene-azaindole derivatives and pyridyl-quinolones derivatives, styryl compounds, styryl-substituted pyridyl compounds, , seleoindoles, selenides, tricyclic polyhydroxylic compounds and benzylphosphonic acid compounds.

Among preferred compounds, it is of interest to focus on pyrimidine derivatives such as N-phenyl-2-pyrimidine-amine derivatives (US 5,521,184 and WO 99/03854), indolinone derivatives and pyrrol-substituted indolinones (US 5,792,783, EP 934 931, US 5,834,504), US 5,883,116, US 5,883,113, US 5, 886,020, WO 96/40116 and WO 00/38519), as well as bis monocyclic, bicyclic aryl and heteroaryl compounds (EP 584 222, US 5,656,643 and WO 92/20642), quinazoline derivatives (EP 602 851, EP 520 722, US 3,772,295 and US 4,343,940), 4-amino-substituted quinazolines (US 3,470,182), 4-thienyl-2-(1H)-quinazolones, 6,7-dialkoxyquinazolines (US 3,800,039), aryl and heteroaryl quinazoline (US 5,721,237, US 5,714,493, US 5,710,158 and WO 95/15758), 4-anilinoquinazoline compounds (US 4,464,375), and 4-thienyl-2-(1H)-quinazolones (US 3,551,427).


So, preferably, the invention relates to a method for treating interstitial cystitis comprising administering a non toxic, potent and selective c-kit inhibitor which is a pyrimidine derivative, more particularly N-phenyl-2-pyrimidine-amine derivatives of formula I :



wherein the R1, R2, R3, R13 to R17 groups have the meanings depicted in EP 564 409 B1, incorporated herein in the description.

20

Preferably, the N-phenyl-2-pyrimidine-amine derivative is selected from the compounds corresponding to formula II :




Wherein R1, R2 and R3 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl or a cyclic or heterocyclic group, especially a pyridyl group;

- 5 R4, R5 and R6 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl, especially a methyl group;

and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least one basic site, such as an amino function.

Preferably, R7 is the following group :



Among these compounds, the preferred are defined as follows:


R1 is a heterocyclic group, especially a pyridyl group,

R2 and R3 are H,

- 15 R4 is a C1-C3 alkyl, especially a methyl group,


R5 and R6 are H,

and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least one basic site, such as an amino function, for example the group :



Therefore, in a preferred embodiment, the invention relates to a method for treating interstitial cystitis comprising the administration of an effective amount of the compound known in the art as CGP57148B :

4-(4-méthylpipérazine-1-ylméthyl)-N-[4-méthyl-3-(4-pyridine-3-yl)pyrimidine-2
 5 ylamino)phényl]-benzamide corresponding to the following formula :



The preparation of this compound is described in example 21 of EP 564 409 and the β -form, which is particularly useful is described in WO 99/03854.

10

Alternatively, the c-kit inhibitor can be selected from :

- indolinone derivatives, more particularly pyrrol-substituted indolinones,
- monocyclic, bicyclic aryl and heteroaryl compounds, quinazoline derivatives,
- and quinaxolines, such as 2-phényl-quinaxoline derivatives, for example 2-phenyl-
 15 6,7-dimethoxy quinaxoline.

In a preferred aspect, the invention contemplates the method mentioned above, wherein said c-kit inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

20

In another embodiment, c-kit inhibitors as mentioned above are inhibitors of activated c-kit. In frame with the invention, the expression "activated c-kit" means a constitutively activated-mutant c-kit including at least one mutation selected from point mutations, deletions, insertions, but also modifications and alterations of the natural c-kit sequence (SEQ ID N°1). Such mutations, deletions, insertions, modifications and alterations can occur in the transphosphorylase domain, in the juxtamembrane domain as well as in any domain directly or indirectly responsible for c-kit activity. The expression "activated c-kit" also means herein SCF-activated c-kit. Preferred and optimal SCF concentrations for activating c-kit are comprised between 5.10^{-7} M and 5.10^{-6} M, preferably around 2.10^{-6} M. In a preferred embodiment, the activated-mutant c-kit in step a) has at least one mutation proximal to Y823, more particularly between amino acids 800 to 850 of SEQ ID No1 involved in c-kit autophosphorylation, notably the D816V, D816Y, D816F and D820G mutants. In another preferred embodiment, the activated-mutant c-kit in step a) has a deletion in the juxtamembrane domain of c-kit. Such a deletion is for example between codon 573 and 579 called c-kit d(573-579). The point mutation V559G proximal to the juxtamembrane domain c-kit is also of interest.

In this regard, the invention contemplates a method for treating interstitial cystitis comprising administering to a human in need of such treatment a compound that is a selective, potent and non toxic inhibitor of activated c-kit obtainable by a screening method which comprises :

- a) bringing into contact (i) activated c-kit and (ii) at least one compound to be tested; under conditions allowing the components (i) and (ii) to form a complex,
- b) selecting compounds that inhibit activated c-kit,
- c) testing and selecting a subset of compounds identified in step b), which are unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

This screening method can further comprise the step consisting of testing and selecting a subset of compounds identified in step b) that are inhibitors of mutant activated c-kit (for example in the transphosphorylase domain), which are also capable of inhibiting SCF-activated c-kit wild.

5 Alternatively, in step a) activated c-kit is SCF-activated c-kit wild.

A best mode for practicing this method consists of testing putative inhibitors at a concentration above 10 μ M in step a). Relevant concentrations are for example 10, 15, 20, 25, 30, 35 or 40 μ M.

10

In step c), IL-3 is preferably present in the culture media of IL-3 dependent cells at a concentration comprised between 0.5 and 10 ng/ml, preferably between 1 to 5 ng/ml.

Examples of IL-3 dependent cells include but are not limited to :

15 - cell lines naturally expressing and depending on c-kit for growth and survival. Among such cells, human mast cell lines can be established using the following procedures :
normal human mast cells can be infected by retroviral vectors containing sequences coding for a mutant c-kit comprising the c-kit signal peptide and a TAG sequence allowing to differentiate mutant c-kits from c-kit wild expressed in hematopoietic cells by
20 means of antibodies.

This technique is advantageous because it does not induce cellular mortality and the genetic transfer is stable and gives satisfactory yields (around 20 %). Pure normal human mast cells can be routinely obtained by culturing precursor cells originating from blood obtained from human umbilical vein. In this regard, heparinated blood from umbilical
25 vein is centrifuged on a Ficoll gradient so as to isolate mononucleated cells from other blood components. CD34+ precursor cells are then purified from the isolated cells mentioned above using the immunomagnetic selection system MACS (Miltenyi biotech).

CD34+ cells are then cultured at 37°C in 5 % CO₂ atmosphere at a concentration of 10⁵ cells per ml in the medium MCCM (α-MEM supplemented with L-glutamine, penicillin, streptomycin, 5 10⁻⁵ M β-mercaptoethanol, 20 % veal foetal serum, 1 % bovine albumin serum and 100 ng/ml recombinant human SCF. The medium is changed every 5 to 7 days. The percentage of mast cells present in the culture is assessed each week, using May-Grünwal Giemsa or Toluidine blue coloration. Anti-tryptase antibodies can also be used to detect mast cells in culture. After 10 weeks of culture, a pure cellular population of mast cells (< 98 %) is obtained.

It is possible using standard procedures to prepare vectors expressing c-kit for transfecting the cell lines established as mentioned above. The cDNA of human c-kit has been described in Yarden et al., (1987) EMBO J.6 (11), 3341-3351. The coding part of c-kit (3000 bp) can be amplified by PCR and cloned, using the following oligonucleotides :

- 5'AAGAAGAGATGGTACCTCGAGGGGTGACCC3' (SEQ ID No2) sens
- 15 - 5'CTGCTTCGCGGCCGCGTTAACTCTTCTCAACCA3' (SEQ ID No3) antisens

The PCR products, digested with NotI and XhoI, has been inserted using T4 ligase in the pFlag-CMV vector (SIGMA), which vector is digested with NotI and XhoI and dephosphorylated using CIP (Biolabs). The pFlag-CMV-c-kit is used to transform bacterial clone XL1-blue. The transformation of clones is verified using the following primers :

- 5'AGCTCGTTTAGTGAACCGTC3' (SEQ ID No4) sens,
- 5'GTCAGACAAAATGATGCAAC3' (SEQ ID No5) antisens.

Directed mutagenesis is performed using relevant cassettes is performed with routine and common procedure known in the art..

The vector Migr-1 (ABC) can be used as a basis for constructing retroviral vectors used for transfecting mature mast cells. This vector is advantageous because it contains the sequence coding for GFP at the 3' and of an IRES. These features allow to select cells infected by the retrovirus using direct analysis with a fluorocytometer. As mentioned
5 above, the N-terminal sequence of c-kit c-DNA can be modified so as to introduce a Flag sequence that will be useful to discriminating heterogeneous from endogenous c-kit.

Other IL-3 dependent cell lines that can be used include but are not limited to:

- BaF3 mouse cells expressing wild-type or mutated form of c-kit (in the
10 juxtamembrane and in the catalytic sites) are described in Kitayama et al, (1996), Blood 88, 995-1004 and Tsujimura et al, (1999), Blood 93, 1319-1329.
- IC-2 mouse cells expressing either c-kit^{WT} or c-kit^{D814Y} are presented in Piao et al, (1996), Proc. Natl. Acad. Sci. USA 93, 14665-14669.

15 IL-3 independent cell lines are :

- HMC-1, a factor-independent cell line derived from a patient with mast cell leukemia, expresses a juxtamembrane mutant c-kit polypeptide that has constitutive kinase activity (Furitsu T et al, J Clin Invest. 1993;92:1736-1744 ; Butterfield et al, Establishment of an immature mast cell line from a patient with mast cell leukemia. Leuk Res. 1988;12:345-
20 355 and Nagata et al, Proc Natl Acad Sci U S A. 1995;92:10560-10564).
- P815 cell line (mastocytoma naturally expressing c-kit mutation at the 814 position) has been described in Tsujimura et al, (1994), Blood 83, 2619-2626.

25 The extent to which component (ii) inhibits activated c-kit can be measured *in vitro* or *in vivo*. In case it is measured *in vivo*, cell lines expressing an activated-mutant c-kit, which

has at least one mutation proximal to Y823, more particularly between amino acids 800 to 850 of SEQ ID No1 involved in c-kit autophosphorylation, notably the D816V, D816Y, D816F and D820G mutants, are preferred.

Example of cell lines expressing an activated-mutant c-kit are as mentioned above.

5

In another preferred embodiment, the method further comprises the step consisting of testing and selecting compounds capable of inhibiting c-kit wild at concentration below 1 μ M. This can be measured *in vitro* or *in vivo*.

Therefore, compounds are identified and selected according to the method described
10 above are potent, selective and non-toxic c-kit wild inhibitors.

Alternatively, the screening method according to the invention can be practiced *in vitro*. In this regard, the inhibition of mutant-activated c-kit and/or c-kit wild can be measured using standard biochemical techniques such as immunoprecipitation and western blot.

15 Preferably, the amount of c-kit phosphorylation is measured.

In a still further embodiment, the invention contemplates a method for treating interstitial cystitis as depicted above wherein the screening comprises :

- a) performing a proliferation assay with cells expressing a mutant c-kit (for example in
20 the transphosphorylase domain), which mutant is a permanent activated c-kit, with a plurality of test compounds to identify a subset of candidate compounds targeting activated c-kit, each having an $IC_{50} < 10 \mu$ M, by measuring the extent of cell death,
- b) performing a proliferation assay with cells expressing c-kit wild said subset of candidate compounds identified in step (a), said cells being IL-3 dependent cells cultured
25 in presence of IL-3, to identify a subset of candidate compounds targeting specifically c-kit,
- c) performing a proliferation assay with cells expressing c-kit, with the subset of compounds identified in step b) and selecting a subset of candidate compounds targeting

c-kit wild, each having an $IC_{50} < 10 \mu M$, preferably an $IC_{50} < 1 \mu M$, by measuring the extent of cell death.

Here, the extent of cell death can be measured by 3H thymidine incorporation, the trypan
5 blue exclusion method or flow cytometry with propidium iodide. These are common techniques routinely practiced in the art.

Therefore, the invention embraces the use of the compounds defined above to manufacture a medicament for treating interstitial cystitis in human.

10 The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

15 In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack
20 Publishing Co., Easton, Pa.).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated
25 as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-
5 cellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

10 Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active
15 compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders,
20 such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

25 Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such

as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Non-lipid polycationic amino
5 polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

The pharmaceutical composition may be provided as a salt and can be formed with many
10 acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succine, acids, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5,
15 that is combined with buffer prior to use.

Pharmaceutical compositions suitable for use in the invention include compositions wherein c-kit inhibitors are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those
20 skilled in the art. A therapeutically effective dose refers to that amount of active ingredient, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio of toxic
25 to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. As mentioned above, a tyrosine kinase inhibitor and more particularly a c-kit

inhibitor according to the invention is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

CLAIMS

1. A method for treating interstitial cystitis comprising administering a tyrosine kinase
5 inhibitor to a human in need of such treatment, wherein said inhibitor is unable to
promote death of IL-3 dependent cells cultured in presence of IL-3.

2. A method according to claim 1, wherein said tyrosine kinase inhibitor is a non-toxic,
selective and potent c-kit inhibitor.

10


3. A method according to claim 2, wherein said inhibitor is selected from the group
consisting of indolinones, pyrimidine derivatives, pyrrolopyrimidine derivatives,
quinazoline derivatives, quinoxaline derivatives, pyrazoles derivatives, bis monocyclic,
bicyclic or heterocyclic aryl compounds, vinylene-azaindole derivatives and pyridyl-
15 quinolones derivatives, styryl compounds, styryl-substituted pyridyl compounds, ,
seleoindoles, selenides, tricyclic polyhydroxylic compounds and benzylphosphonic acid
compounds.

4. A method for treating interstitial cystitis comprising administering a non toxic, potent
20 and selective c-kit inhibitor to a human in need of such treatment, selected from the
group consisting of :

- pyrimidine derivatives, more particularly N-phenyl-2-pyrimidine-amine derivatives.
- indolinone derivatives, more particularly pyrrol-substituted indolinones,
- monocyclic, bicyclic aryl and heteroaryl compounds,
- 25 - and quinazoline derivatives,

wherein said inhibitor is unable to promote death of IL-3 dependent cells cultured in
presence of IL-3.

5. A method according to claim 2, wherein said inhibitor is selected from the group consisting of N-phenyl-2-pyrimidine-amine derivatives having the formula II :




5

Wherein R1, R2 and R3 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl or a cyclic or heterocyclic group, especially a pyridyl group;

R4, R5 and R6 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl, especially a methyl group;

10

and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least one basic site, such as an amino function, preferably the following group :



15 6. A method according to claim 5, wherein said inhibitor is the 4-(4-méthylpipérazine-1-ylméthyl)-N-[4-méthyl-3-(4-pyridine-3-yl)pyrimidine-2 ylamino]phényl]-benzamide.

7. A method according to claim 5 or 6, wherein said c-kit inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

20

8. A method according to claim 2, wherein said inhibitor is an inhibitor of activated c-kit selected from a constitutively activated-mutant c-kit and/or SCF-activated c-kit.
9. A method according to claim 8, wherein the activated-mutant c-kit has at least one
5 mutation selected from mutations proximal to Y823, more particularly between amino acids 800 to 850 of SEQ ID No1 involved in c-kit autophosphorylation, notably the D816V, D816Y, D816F and D820G mutants, and a deletion in the juxtamembrane domain of c-kit, preferably between codon 573 and 579.
- 10 10. A method for treating interstitial cystitis comprising administering to a human in need of such treatment a compound that is a selective, potent and non toxic inhibitor of activated c-kit obtainable by a screening method which comprises :
- a) bringing into contact (i) activated c-kit and (ii) at least one compound to be tested; under conditions allowing the components (i) and (ii) to form a complex,
- 15 b) selecting compounds that inhibit activated c-kit,
- c) testing and selecting a subset of compounds identified in step b), which are unable to promote death of IL-3 dependent cells cultured in presence of IL-3.
11. A method according to claim 10, wherein the screening method further comprises the
20 step consisting of testing and selecting a subset of compounds identified in step b) that are inhibitors of mutant activated c-kit, which are also capable of inhibiting SCF-activated c-kit wild.
12. A method according to claim 10, wherein activated c-kit is SCF-activated c-kit wild.
- 25 13. A method according to one of claims 10 to 12, wherein putative inhibitors are tested at a concentration above 10 μ M in step a).

14. A method according to one of claims 10 to 13, wherein IL-3 is present in the culture media of IL-3 dependent cells at a concentration comprised between 0.5 and 10 ng/ml, preferably between 1 to 5 ng/ml.

5 15. A method according to one of claims 10 to 14, wherein the extent to which component (ii) inhibits activated c-kit can be measured *in vitro* or *in vivo*.

16. A method according to one of claims 10 to 15 wherein, the screening method further comprises the step consisting of testing and selecting *in vitro* or *in vivo* compounds
10 capable of inhibiting c-kit wild at concentration below 1 μ M.

17. A method according to one of claims 10 to 16 wherein, the test is performed using cells lines selected from the group consisting of mast cells, transfected mast cells, BaF3, and IC-2.
15

18. A method according to one of claims 10 to 17 wherein, the test includes the determination of the amount of c-kit phosphorylation.

19. A method for treating interstitial cystitis according to one of claims 10 to 17, wherein
20 the screening comprises :

- a) performing a proliferation assay with cells expressing a mutant c-kit (for example in the transphosphorylase domain), which mutant is a permanent activated c-kit, with a plurality of test compounds to identify a subset of candidate compounds targeting activated c-kit, each having an $IC_{50} < 10 \mu$ M, by measuring the extent of cell death,
- 25 b) performing a proliferation assay with cells expressing c-kit wild said subset of candidate compounds identified in step (a), said cells being IL-3 dependent cells cultured in presence of IL-3, to identify a subset of candidate compounds targeting specifically c-kit,

c) performing a proliferation assay with cells expressing c-kit, with the subset of compounds identified in step b) and selecting a subset of candidate compounds targeting c-kit wild, each having an $IC_{50} < 10 \mu M$, preferably an $IC_{50} < 1 \mu M$, by measuring the extent of cell death.

5

20. A method according to one of claims 1 to 19, wherein the inhibitor is administered orally.

21. Use of a tyrosine kinase inhibitor, more particularly a c-kit inhibitor, to manufacture
10 a medicament for treating interstitial cystitis in human.

22. Use according to claim 21, wherein said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

SEQUENCE LISTING

<110> AB Science

<120> Use of potent, selective and non toxic c-kit inhibitors
for treating interstitial cystitis

<130> D19832 NT

<150> US 60/323,315

<151> 2001-09-20

<160> 5

<170> PatentIn Ver. 2.1

<210> 1

<211> 976

<212> PRT

<213> Homo sapiens

<220>

<223> Human c-kit

<400> 1

Met Arg Gly Ala Arg Gly Ala Trp Asp Phe Leu Cys Val Leu Leu Leu
1 5 10 15

Leu Leu Arg Val Gln Thr Gly Ser Ser Gln Pro Ser Val Ser Pro Gly
20 25 30

Glu Pro Ser Pro Pro Ser Ile His Pro Gly Lys Ser Asp Leu Ile Val
35 40 45

Arg Val Gly Asp Glu Ile Arg Leu Leu Cys Thr Asp Pro Gly Phe Val
50 55 60

Lys Trp Thr Phe Glu Ile Leu Asp Glu Thr Asn Glu Asn Lys Gln Asn
65 70 75 80

Glu Trp Ile Thr Glu Lys Ala Glu Ala Thr Asn Thr Gly Lys Tyr Thr
85 90 95

Cys Thr Asn Lys His Gly Leu Ser Asn Ser Ile Tyr Val Phe Val Arg
100 105 110

Asp Pro Ala Lys Leu Phe Leu Val Asp Arg Ser Leu Tyr Gly Lys Glu
115 120 125

Asp Asn Asp Thr Leu Val Arg Cys Pro Leu Thr Asp Pro Glu Val Thr
130 135 140

Asn Tyr Ser Leu Lys Gly Cys Gln Gly Lys Pro Leu Pro Lys Asp Leu
145 150 155 160

Arg Phe Ile Pro Asp Pro Lys Ala Gly Ile Met Ile Lys Ser Val Lys
165 170 175

Arg Ala Tyr His Arg Leu Cys Leu His Cys Ser Val Asp Gln Glu Gly
180 185 190

Lys Ser Val Leu Ser Glu Lys Phe Ile Leu Lys Val Arg Pro Ala Phe
195 200 205

Lys Ala Val Pro Val Val Ser Val Ser Lys Ala Ser Tyr Leu Leu Arg
210 215 220

Glu Gly Glu Glu Phe Thr Val Thr Cys Thr Ile Lys Asp Val Ser Ser
 225 230 235 240
 Ser Val Tyr Ser Thr Trp Lys Arg Glu Asn Ser Gln Thr Lys Leu Gln
 245 250 255
 Glu Lys Tyr Asn Ser Trp His His Gly Asp Phe Asn Tyr Glu Arg Gln
 260 265 270
 Ala Thr Leu Thr Ile Ser Ser Ala Arg Val Asn Asp Ser Gly Val Phe
 275 280 285
 Met Cys Tyr Ala Asn Asn Thr Phe Gly Ser Ala Asn Val Thr Thr Thr
 290 295 300
 Leu Glu Val Val Asp Lys Gly Phe Ile Asn Ile Phe Pro Met Ile Asn
 305 310 315 320
 Thr Thr Val Phe Val Asn Asp Gly Glu Asn Val Asp Leu Ile Val Glu
 325 330 335
 Tyr Glu Ala Phe Pro Lys Pro Glu His Gln Gln Trp Ile Tyr Met Asn
 340 345 350
 Arg Thr Phe Thr Asp Lys Trp Glu Asp Tyr Pro Lys Ser Glu Asn Glu
 355 360 365
 Ser Asn Ile Arg Tyr Val Ser Glu Leu His Leu Thr Arg Leu Lys Gly
 370 375 380
 Thr Glu Gly Gly Thr Tyr Thr Phe Leu Val Ser Asn Ser Asp Val Asn
 385 390 395 400
 Ala Ala Ile Ala Phe Asn Val Tyr Val Asn Thr Lys Pro Glu Ile Leu
 405 410 415
 Thr Tyr Asp Arg Leu Val Asn Gly Met Leu Gln Cys Val Ala Ala Gly
 420 425 430
 Phe Pro Glu Pro Thr Ile Asp Trp Tyr Phe Cys Pro Gly Thr Glu Gln
 435 440 445
 Arg Cys Ser Ala Ser Val Leu Pro Val Asp Val Gln Thr Leu Asn Ser
 450 455 460
 Ser Gly Pro Pro Phe Gly Lys Leu Val Val Gln Ser Ser Ile Asp Ser
 465 470 475 480
 Ser Ala Phe Lys His Asn Gly Thr Val Glu Cys Lys Ala Tyr Asn Asp
 485 490 495
 Val Gly Lys Thr Ser Ala Tyr Phe Asn Phe Ala Phe Lys Gly Asn Asn
 500 505 510
 Lys Glu Gln Ile His Pro His Thr Leu Phe Thr Pro Leu Leu Ile Gly
 515 520 525
 Phe Val Ile Val Ala Gly Met Met Cys Ile Ile Val Met Ile Leu Thr
 530 535 540
 Tyr Lys Tyr Leu Gln Lys Pro Met Tyr Glu Val Gln Trp Lys Val Val
 545 550 555 560
 Glu Glu Ile Asn Gly Asn Asn Tyr Val Tyr Ile Asp Pro Thr Gln Leu
 565 570 575

Pro Tyr Asp His Lys Trp Glu Phe Pro Arg Asn Arg Leu Ser Phe Gly
 580 585 590
 Lys Thr Leu Gly Ala Gly Ala Phe Gly Lys Val Val Glu Ala Thr Ala
 595 600 605
 Tyr Gly Leu Ile Lys Ser Asp Ala Ala Met Thr Val Ala Val Lys Met
 610 615 620
 Leu Lys Pro Ser Ala His Leu Thr Glu Arg Glu Ala Leu Met Ser Glu
 625 630 635 640
 Leu Lys Val Leu Ser Tyr Leu Gly Asn His Met Asn Ile Val Asn Leu
 645 650 655
 Leu Gly Ala Cys Thr Ile Gly Gly Pro Thr Leu Val Ile Thr Glu Tyr
 660 665 670
 Cys Cys Tyr Gly Asp Leu Leu Asn Phe Leu Arg Arg Lys Arg Asp Ser
 675 680 685
 Phe Ile Cys Ser Lys Gln Glu Asp His Ala Glu Ala Ala Leu Tyr Lys
 690 695 700
 Asn Leu Leu His Ser Lys Glu Ser Ser Cys Ser Asp Ser Thr Asn Glu
 705 710 715 720
 Tyr Met Asp Met Lys Pro Gly Val Ser Tyr Val Val Pro Thr Lys Ala
 725 730 735
 Asp Lys Arg Arg Ser Val Arg Ile Gly Ser Tyr Ile Glu Arg Asp Val
 740 745 750
 Thr Pro Ala Ile Met Glu Asp Asp Glu Leu Ala Leu Asp Leu Glu Asp
 755 760 765
 Leu Leu Ser Phe Ser Tyr Gln Val Ala Lys Gly Met Ala Phe Leu Ala
 770 775 780
 Ser Lys Asn Cys Ile His Arg Asp Leu Ala Ala Arg Asn Ile Leu Leu
 785 790 795 800
 Thr His Gly Arg Ile Thr Lys Ile Cys Asp Phe Gly Leu Ala Arg Asp
 805 810 815
 Ile Lys Asn Asp Ser Asn Tyr Val Val Lys Gly Asn Ala Arg Leu Pro
 820 825 830
 Val Lys Trp Met Ala Pro Glu Ser Ile Phe Asn Cys Val Tyr Thr Phe
 835 840 845
 Glu Ser Asp Val Trp Ser Tyr Gly Ile Phe Leu Trp Glu Leu Phe Ser
 850 855 860
 Leu Gly Ser Ser Pro Tyr Pro Gly Met Pro Val Asp Ser Lys Phe Tyr
 865 870 875 880
 Lys Met Ile Lys Glu Gly Phe Arg Met Leu Ser Pro Glu His Ala Pro
 885 890 895
 Ala Glu Met Tyr Asp Ile Met Lys Thr Cys Trp Asp Ala Asp Pro Leu
 900 905 910
 Lys Arg Pro Thr Phe Lys Gln Ile Val Gln Leu Ile Glu Lys Gln Ile
 915 920 925

Ser Glu Ser Thr Asn His Ile Tyr Ser Asn Leu Ala Asn Cys Ser Pro
 930 935 940
 Asn Arg Gln Lys Pro Val Val Asp His Ser Val Arg Ile Asn Ser Val
 945 950 955 960
 Gly Ser Thr Ala Ser Ser Ser Gln Pro Leu Leu Val His Asp Asp Val
 965 970 975

<210> 2
 <211> 30
 <212> DNA
 <213> Homo sapiens

<220>
 <223> Primer

<400> 2
 aagaagagat ggtacctcga ggggtgaccc 30

<210> 3
 <211> 33
 <212> DNA
 <213> Homo sapiens

<220>
 <223> Primer

<400> 3
 ctgcttcgcg gccgcgttaa ctcttctcaa cca 33

<210> 4
 <211> 20
 <212> DNA
 <213> Homo sapiens

<220>
 <223> Primer

<400> 4
 agctcgttta gtgaaccgtc 20

<210> 5
 <211> 20
 <212> DNA
 <213> Homo sapiens

<220>
 <223> Primer

<400> 5
 gtcagacaaa atgatgcaac 20